

Note to readers with disabilities: *EHP* strives to ensure that all journal content is accessible to all readers. However, some figures and Supplemental Material published in *EHP* articles may not conform to [508 standards](#) due to the complexity of the information being presented. If you need assistance accessing journal content, please contact ehponline@niehs.nih.gov. Our staff will work with you to assess and meet your accessibility needs within 3 working days.

Supplemental Material

Heat-Related Mortality in Japan after the 2011 Fukushima Disaster: An Analysis of Potential Influence of Reduced Electricity Consumption

Yoonhee Kim, Antonio Gasparri, Masahiro Hashizume, Yasushi Honda, Chris Fook Sheng Ng, and Ben Armstrong

Table of Contents

Table S1. Summary statistics of power use, mortality, and daily mean temperature across 47 prefectures in Japan by less- and most-affected areas

Table S2. Mean cumulative relative risks and ratio of relative risks with 95% confidence intervals in less-affected and most-affected areas for people aged 65 years and over

Table S3. Cumulative relative risks (RRs) and the ratio of relative risk (RRRs) for all-cause mortality associated with heat (defined as the 95th percentile of daily mean temperature during May–September) over 0–10 lag days, compared with those for the 50th percentile of the temperature, combined by meta-analysis across 13 prefectures in the most-affected area and 95% confidence intervals, based on the interrupted time-series (ITS) analysis with and without adjustment for an air pollutant (2-days moving average)

Figure S1. Time-series plots of the daily number of all-cause deaths (A) and daily mean

temperature (B) in Tokyo and daily electricity consumption across the eight prefectures served by the Tokyo Electric Power Company (TEPCO) listed in Table S1. The vertical line indicates the date of the Great East Japan Earthquake occurred on 11 March 2011.

Figure S2. Forest plot for prefecture-specific ratio of relative risks (RRRs) for all-cause mortality associated with heat (the 99th percentile of daily mean temperature during May–September for prefecture specific shown in Table S1) compared with the 50th percentile of the temperature over 0–10 lag days after to before the earthquake and 95% confidence intervals based on the distributed lag non-linear model adjusted for day-of-week or holiday, date, and day-in-season (natural cubic spline with 4 df/year) \times year. The periods for before and after correspond to 2008–2010 and 2011–2012, respectively. A fixed effect meta-analysis was used to estimate the summary RRRs in less-affected and most-affected areas.

Figure S3. Prefecture-specific cumulative relative risks (RRs) for all-cause mortality associated with daily mean temperature during May–September for prefecture specific (the distribution of the temperature shown in Table S1) compared with the 50th percentile of the temperature over 0–10 lag days before and after the earthquake and 95% confidence intervals based on the distributed lag non-linear model adjusted for day-of-week or holiday, date, and day-in-season (natural cubic spline with 4 df/year) \times year (1st column), their lag-response curves of the RRs for the 95th percentile of daily mean temperature (2nd column), and changes in the RRs after to before (ratio of relative risks, RRRs) (3rd column). The shaded areas indicate 95% confidence intervals.

Figure S4. Cumulative relative risks for all-cause (ALL), cardiovascular (CVD), and respiratory (RESP) mortality for people aged 65 and over associated with heat at the 95th (A) or 99th (B) percentiles of daily mean temperature during May–September in Tokyo (30.1°C or 31.1°C, respectively), compared with those for the 50th percentile of the temperature (24.4°C) over 0–10 lag days, stratified by the estimated percent reduction in

electricity consumption (i.e. the difference between observed and expected consumption after the earthquake based on a model adjusted for temperature, day-of-week or holiday, and day-in-season using daily electricity consumption in the TEPCO service area that includes Tokyo).

Figure S5. Results by sensitivity analysis excluding the 2011 data showing forest plot for prefecture-specific ratio of relative risks (RRRs) for all-cause mortality associated with heat (the 95th percentile of daily mean temperature during May–September for prefecture specific shown in Table S1) compared with the 50th percentile of the temperature over 0–10 lag days after to before the earthquake and 95% confidence intervals based on the distributed lag non-linear model adjusted for day-of-week or holiday, date, and day-in-season (natural cubic spline with 4 df/year) \times year. The periods for before and after correspond to 2008–2010 and 2011–2012, respectively. A fixed effect meta-analysis was used to estimate the summary RRRs in less-affected and most-affected areas. The results were consistent with the main results although the 95% confidence intervals became wider due to the smaller statistical power.

Figure S6. Results by sensitivity analyses. Estimated ratio of relative risks (RRRs) with 95% confidence intervals (CIs) in Tokyo based on the interrupted time-series analysis, according to increases in (A) degree of piecewise polynomials of B-spline basis for temperature, (B) degree of freedom of the natural cubic spline for day-in-season, (C) the place of knot for the spline of temperature, and (D) the placement of reference temperature. (E) The cumulative relative risks for all-cause mortality associated with daily mean temperature during May–September compared with the 50th percentile of the temperature over 0–21 lag days and (F) their lag-response curves for the 95th percentile of daily mean temperature in Tokyo. The shaded areas in panels (E) and (F) indicate 95% confidence intervals. (G) Deviance residuals over time and (H) partial autocorrelation function based on the final model of interrupted time-series analysis in Tokyo.

Figure S7. Results by sensitivity analyses. The cumulative relative risks for all-cause mortality associated with daily mean temperature during May–September compared with the 50th percentile of the temperature over 0–10 lag days in Tokyo based on the interrupted time-series analysis, applying a range of number of knots for the quadratic B-spline of daily mean temperature from 1 knot to 4 knots. The shaded areas in each panel indicate 95% confidence intervals.